

First Russian record of *Erpobdella monostrata*: DNA barcoding and geographical distribution

(Annelida, Hirudinida, Erpobdellidae)

Serge Utevsky, Pavel G. Dubov & Alexander A. Prokin

Utevsky, S., Dubov, P. G. & Prokin, A. A. 2015. First Russian record of *Erpobdella monostrata*: DNA barcoding and geographical distribution (Annelida, Hirudinida, Erpobdellidae). *Spixiana* 38(2): 161-168.

New information on the range of *Erpobdella monostrata* (Lindenfeld & Pietruszynski, 1890) was obtained. Results of our study show that this leech occurs not only in Central Europe but reaches European Russia. However, *E. monostrata* has not been recorded from the vast interspace between its localities in Central Europe and a dystrophic lake in the Voronezh Oblast in Russia, the easternmost spot of its range known so far. The taxonomic identity of the leeches was confirmed both by a morphological examination and a molecular phylogenetic analysis based on COI sequences. Genetic differentiation between western and eastern samples of the species points towards exploring more specimens and markers. If not disjunct, the present geographical gap of records may be explained by low population sizes and the scarcity of suitable habitats; recent climate changes might cause population expansions in the east.

Serge Utevsky, Department of Zoology and Animal Ecology, V. N. Karazin Kharkiv National University, Maidan Svobody 4, Kharkiv 61022, Ukraine; e-mail: sutevsk@univer.kharkov.ua

Pavel G. Dubov, Voronezh State University, Universitetskaya sq., 1, Voronezh 394006, Russia

Alexander A. Prokin, Voronezh State University, Universitetskaya sq., 1, Voronezh 394006, Russia; and Papanin Institute for Biology of Inland Waters, Russian Academy of Sciences, Borok, Nekouz district, Yaroslavl Oblast 152742, Russia; e-mail: prokina@mail.ru

Introduction

Erpobdellid leeches are common members of fresh-water benthic communities. They are macrophagous predators of aquatic invertebrates originating from hematophagous ancestors (Siddall 2002). Some of the species in this group have been used in ecological studies and water quality assessment (Lukin 1976, Schenková et al. 2007, Neesemann & Moog 2002). The classical systematics of erpobdellids was based on the annulation of segments. In the molecular phylogenetic era, the importance of annulation has been reconsidered and the genus *Erpobdella* sensu stricto

has been found non-monophyletic (Siddall 2002). Further phylogenetic analyses and taxonomic revisions are needed to reveal the evolutionary history of erpobdellids and resolve taxonomic uncertainties.

The taxonomy of a number of common erpobdellid species has been confused or neglected. The taxonomic identity of *Erpobdella monostrata* (Lindenfeld & Pietruszynski, 1890) was confused for a long time since its formal description. This taxon was initially described as *Nepheleis octoculata* var. *monostrata* Lindenfeld & Pietruszynski, 1890. Then the preoccupied epithet “*monostrata*” was employed again to designate another leech taxon, *Herpobdella*

atomaria var. *monostriata* Gedroyć, 1916, which is currently recognized as a synonym of the correct designation of this species, *Erpobdella vilnensis* (Liskiewicz, 1925) (Agapow & Bielecki 1992, Neseemann & Neubert 1999).

In his seminal monographs, Lukin (1962, 1976) mentioned only "*Erpobdella monostriata*" (Gedroyć, 1916) for the Carpathians and adjacent areas and did not record *Erpobdella monostriata* (Lindenfeld & Pietruszynski, 1890) for Ukraine and the former USSR. Recently Utevsky et al. (2012) have recorded *Erpobdella vilnensis* (Liskiewicz, 1925) from the Ukrainian Carpathians and south-western Ukraine and by doing so fixed the nomenclatural problem of confusing *E. vilnensis* with *E. monostriata* at the putative eastern border of the range of the former. At the same time, the geographical distribution of *E. monostriata* is still poorly studied. It has been unclear how far its range extends to the east. Until recently *Erpobdella monostriata* was not found in the former Soviet Union (Lukin 1962, 1976) and newly independent countries of Eastern Europe. This research addresses the information gap in our knowledge of the geographical distribution of *E. monostriata*.

Material and methods

Sample collection

Samples of *E. monostriata* were collected during the summer season of 2011 and on 16th June, 2013 from a small forest lake Ugolnoe in the Usman Pine Forest in the Voronezh Oblast, Russia near the village of Maklok (51°48'40.20" N, 39°24'23.58" E), relaxed in 10 % ethanol, fixed and preserved in 96 % ethanol. Altogether, 37 individuals were studied by using morphological and molecular methods. A newly sequenced specimen of *Erpobdella vilnensis* from the Mykolayiv Oblast, Ukraine (see Utevsky et al. 2012) was also used for the analysis. The specimens are stored in the collection of invertebrates of the Department of Zoology and Animal Ecology, V. N. Karazin Kharkiv National University.

Measurements

Measurements were made with an eyepiece micrometer. Arithmetical means of the body length and width and their standard errors were calculated. The measurements were based on 32 specimens.

Mapping

Geographical coordinates were assigned to the new and previous localities where *E. monostriata* was found. Agapow & Bielecki (1992) and later publications were considered. In the field, the geographical position was determined using a global positioning system (GPS) device. In other cases, localities were identified according to the description in the literature.

The Google Earth (version 4.3) online application (<http://earth.google.com/>) was used. The record points were plotted on a map using QGIS 2.0.

DNA extraction, amplification and sequencing

Small pieces (approx. 5×2×1 mm) of skin and muscle tissue were cut from the lateral part of the body. Care was taken not to reach the digestive system, which often contains remnants of unknown prey species. Genomic DNA was isolated using the GENE ELUTE Mammalian Genomic DNA minprep kit from Sigma-Aldrich (Steinheim, Germany).

The mitochondrial cytochrome oxidase subunit one (COI) fragment was amplified using LSO and HCO primers described in Folmer et al. (1994). PCR was performed by applying 35 cycles of 1 min at 94 °C, 1 min at 46 °C, and 2.5 min at 72 °C, following a 4 min denaturation step at 94 °C. In order to purify the PCR products two enzymes, Exonuclease I and Shrimp alkaline phosphatase (SAP) (Fermentas, Thermo Fisher Scientific, USA), were used. Exonuclease I (0.2 µl) and SAP (1 µl) were added to 10 µl of the PCR product. After that, the mixture was incubated for 45 min at 37 °C and followed by a 15 min incubation at 80 °C. The purified products were sequenced in both directions with amplification primers under BIG DYE Terminator cycling conditions, purified by ethanol precipitation, and run on an Applied Biosystems 3730xl sequencer by Macrogen (Seoul, Korea). Sequence chromatograms were edited and assembled with the help of ChromasPro 1.32 (Technelysium Pty., Queensland, Australia). The correctness of COI sequences was verified at the amino acid level. The obtained sequences were assigned GenBank Accession Numbers. Some COI sequences belonging to other erpobdellid species from previous studies were also used for inferring a phylogenetic tree (Table 1).

Phylogenetic analysis

The COI sequences were unambiguously aligned using ClustalW associated with BioEdit v7.1.7 (Hall 1999). The length of the aligned COI sequences was 660 bp. The best-fit models for each partition, the first, second and third codon positions of COI, were identified with the Bayesian information criterion (Schwarz 1978) using KAKUSAN4 (Tanabe 2011): GTR with proportion of invariant sites (+I), HKY85+I and HKY85 with gamma distribution (+G) respectively.

Phylogenetic relationships were determined by Bayesian inference using MrBayes v3.1.2. (Ronquist & Huelsenbeck 2003). Two arhynchobdellid species *Hirudo orientalis* Utevsky et Trontelj, 2005 (Hirudiniformes) and *Orobdella jimai* Oka, 1895 (Erpobdelliformes) were used as outgroup taxa to reveal the phylogenetic relationships between erpobdellid taxa. Searches were performed in two parallel runs with four chains each for five million generations, sampled every 100th generation. After discarding the first 25 % of the sampled trees, final topologies were consented following the 50 %

majority rule. Numbers of base differences per site (*p*-distances) and their standard errors were calculated using MEGA5 (Tamura et al. 2011).

Habitat

The basin of the Ugolnoe Lake is a former riverbed of a left-bank tributary of the Voronezh River, ceased to exist during the formation of the estuary of the Usman River during an activation of the Voronezh neotectonic uplift (Khlyzova et al. 2007). The name “ugolnoe” comes from a facility that was located here in the late 19th and early 20th centuries to produce birch charcoal. According to hydrochemical characteristics determined by Zhi-votova & Koroteeva (2002), the lake should be classified as a dystrophic (humic) water.

Forest wildfires of 2010 had a significant impact on the hydrochemical and hydrobiological regime of waters of the Usman Pine Forest, including the Ugolnoe Lake. It is known that after wildfires the absolute and relative (compared to nitrogen) content of organic carbon and a number of ions increased in surface waters due to the accumulation of ash and fly ash from the

catchment area; the balance of biogeochemical fluxes changes due to dramatic alterations in vegetation and soil permeability (Bitner et al. 2001, Meyer et al. 1995, Strauss & Lamberti 2000).

Results

Morphology

External characters (Fig. 1) of the specimens examined are in agreement with the previous description of *E. monostrata* by Nesemann & Neubert (1999). The leeches are small, the average total length including suckers is 15.13 ± 0.56 mm (range 8.50–21.00 mm). The maximum body width is 2.16 ± 0.12 mm (range 1.50–5.20 mm). The body surface is smooth, papillae are lacking. A mid-body segment consists of five annuli equal in length. The gonopores are separated by four annuli. The coloration of living leeches is light brownish. All preserved specimens have a mid-dorsal dark longitudinal stripe located along

Table 1. Collection sites and sequence accession data for the leech species analysed.

Species	Collection site	GenBank accession No.	Reference
<i>Erpobdella monostrata</i> (Lindenfeld & Pietruszynski, 1890)	Voronezh Oblast, Russia	KP300764	This study
<i>Erpobdella monostrata</i> (Lindenfeld & Pietruszynski, 1890)	Mecklenburg-Western Pomerania, Lake Neustädter See, Germany	HM246601	Trajanovski et al. 2010
<i>Erpobdella monostrata</i> (Lindenfeld & Pietruszynski, 1890)	Magdeburg, Germany	DQ009665	Pfeiffer et al. 2005
<i>Erpobdella nigricollis</i> (Brandes, 1899)	Kassel-Wehlheiden, Germany	DQ009664	Pfeiffer et al. 2005
<i>Erpobdella nigricollis</i> (Brandes, 1899)	Mecklenburg-Western Pomerania, Lake Neustädter See, Germany	HM246603	Trajanovski et al. 2010
<i>Erpobdella testacea</i> (Savigny, 1820)	France	AF116027	Apakupakul et al. 1999
<i>Erpobdella vilnensis</i> (Liskiewicz, 1925)	A stream flowing to the Southern Bug, Mykolayiv Oblast, Ukraine	KP300763	This study, Utevsky et al. 2012
<i>Erpobdella vilnensis</i> (Liskiewicz, 1925)	Wörlitzer Park, Germany	DQ009663	Pfeiffer et al. 2005
<i>Erpobdella vilnensis</i> (Liskiewicz, 1925)	Mecklenburg-Western Pomerania, Germany	HM246585	Trajanovski et al. 2010
<i>Erpobdella octoculata</i> (Linnaeus, 1758)	Uzbekistan	HQ336344	Oceguera-Figueroa et al. 2011
<i>Erpobdella octoculata</i> (Linnaeus, 1758)	Lake Ohrid, Macedonia	HM246555	Trajanovski et al. 2010
<i>Erpobdella octoculata</i> (Linnaeus, 1758)	Lake Prespa, Macedonia	HM246599	Trajanovski et al. 2010
<i>Dina ohridana</i> Sket, 1968	Lake Ohrid, Albania/Macedonia,	HM246594	Trajanovski et al. 2010
<i>Dina lepinja</i> Sket & Šapkarev, 1986	Macedonia, Lake Ohrid	HM246539	Trajanovski et al. 2010
<i>Dina lineata</i> (O.F. Müller, 1774)	Mecklenburg-Western Pomerania, Germany	HM246611	Trajanovski et al. 2010
<i>Trocheta haskonis</i> Grosser, 2000	Wörlitzer Park, Germany	DQ009668	Pfeiffer et al. 2005
<i>Orobdella jimai</i> Oka, 1895	Nagano, Japan	AB679674	Nakano 2012
<i>Hirudo orientalis</i> Utevsky & Trontelj, 2005	Azerbaijan	JN104645	Trontelj & Utevsky 2012




Fig. 1. *Erpobdella monostrata* (Lindenfeld & Pietruszynski, 1890). **A.** Entire specimen with a characteristic mid-dorsal dark longitudinal stripe. **B.** Clitellum, ventral view, with two gonopores separated by four annuli: M, male gonopore; F, female gonopore.

the entire body without interruption. The stripe is blurred and associated with clouds of dark pigment in many specimens. In most of living leeches, the stripe is indistinct or lacking whereas this distinguishing feature emerges in specimens fixed and preserved in strong ethanol.

Barcoding and phylogeny

The resulting Bayesian tree (Fig. 2) indicates that the sample of the leech specimens collected in Russia joined a clade with two other samples of *E. monostrata* from Germany. The posterior probability of the monophyletic group is 1.00. The *p*-distances between the Russian sample and two sequences HM246601 and DQ009665 are 0.015 ± 0.005 and 0.018 ± 0.006 respectively. The *p*-distance between the two samples originating from Germany is as low as 0.004 ± 0.003 . The mean *p*-distance within *E. monostrata* was 0.014 ± 0.005 (Table 2). The distances between the




Fig. 2. Phylogenetic relationships between major groups of Palaeartic erpobdellids obtained by Bayesian inference and based on COI sequences. Posterior probabilities are shown for clades. The tree is rooted at *Hirudo orientalis*.

Erpobdella species are shown in Table 3. The sample of *E. vilnensis* collected in south-western Ukraine (Utevsky et al. 2012) joined a well-supported clade with its conspecifics.

Geographical distribution

Altogether 64 localities of *E. monostrata* were considered. Its range reaches as far to the west as $06^{\circ}22'07.82''$ in the Netherlands. The easternmost locality is $39^{\circ}24'23.58''$, a new record from the Russian Federation. The northern frontier of the explored

Table 2. The number of base differences per site from averaging over all sequence pairs within each *Erpobdella* species (π) and its standard error (based on *p*-distances).

Species	π
<i>E. monostrata</i>	0.014 ± 0.005
<i>E. nigricollis</i>	0.002 ± 0.001
<i>E. vilnensis</i>	0.018 ± 0.006
<i>E. octoculata</i>	0.057 ± 0.022



Fig. 3. Geographical distribution of *Erpobdella monostrata* (Lindenfeld & Pietruszynski, 1890). Previous records are mapped with circles and the new record from Russia is represented by a star.

range of *E. monostrata* is 56°04'08.73" in Sweden (Fig. 3). The southern frontier of the range is still unclear (see below). Reliable records of *E. monostrata* were found for eight European countries: the Netherlands (Haaren et al. 2004), Germany (Grosser 1996, 2003, Jueg 1998, 2013, Jueg et al. 2007, Haesloop 2002, Pfeiffer et al. 2005, Westendorff et al. 2008, Trajanovski et al. 2010, Enting & Arndt-Dietrich 2012), Denmark (Trontelj et al. 1996), Sweden (Uwe Jueg, personal communication), Poland (Agapow & Bielecki 1992, Fleituch 2003, Bielecki et al. 2004, 2011a,b, Szczesny 2005, Nabdodnik et al. 2004, Agapow et al. 2006, Agapow & Nabdodnik 2006, Agapow

et al. 2008, Raczynska & Chojnacki 2009, Koperski 2010, Koszalka 2012), Lithuania (Uwe Jueg, personal communication), Belarus (Haaren et al. 2004) and Russia (this research).

Discussion

Both morphological and molecular analyses showed that the leeches sampled in the Ugolnoe Lake, Voronezh Oblast, Russia should be assigned to *Erpobdella monostrata* (Lindenfeld & Pietruszynski, 1890). The leeches possess a distinguishing feature of the spe-

Table 3. The number of base differences per site from averaging over all sequence pairs between *Erpobdella* species and their standard errors (based on *p*-distances).

Species	1	2	3	4
1 <i>E. monostrata</i>				
2 <i>E. nigricollis</i>	0.096±0.012			
3 <i>E. testacea</i>	0.131±0.013	0.116±0.012		
4 <i>E. vilnensis</i>	0.117±0.013	0.107±0.013	0.132±0.013	
5 <i>E. octoculata</i>	0.139±0.013	0.117±0.012	0.124±0.012	0.123±0.012

cies, the mid-dorsal dark longitudinal stripe located along the entire body. That coloration pattern has never been found in other *Erpobdella* species. The COI tree recovers a well-supported clade of *Erpobdella monostrata*. The eastern specimen differs by less than 2% from GenBank sequences of two western *E. monostrata*, but the latter form a strongly supported clade, suggesting geographic structure. Being more than eight times higher than within species differences of *E. monostrata*, the *p*-distance between *E. monostrata* and other *Erpobdella* species ranges from 0.096 to 0.139. This can be considered as reliable evidence of the species status of the *E. monostrata* clade (see Birky 2013). Our analyses also corroborated species-level genetic differences between *E. monostrata* and *E. vilnensis* that went through a long period of nomenclatural confusions (see Agapow & Bielecki 1992). Further analyses dealing with more samples and diverse molecular markers are necessary to reveal accurate intraspecific relationships within *E. monostrata* and its place among other erpobdellid taxa.

Recent studies suggest that ranges of a number of Western Palaearctic leech species extend more easterly than considered before. For example, *Hirudo verbana* Carena, 1820 occurs in Uzbekistan (Utevsky et al. 2010) and *E. vilnensis* has been recorded from south-western Ukraine (Utevsky et al. 2012) and Kyrgyzstan (Jueg et al. 2013). Our research, in turn, indicates that *E. monostrata*, recorded mainly from Central Europe, reaches the Don basin in Russia in the eastern portion of its known range. Records from the Czech Republic, Slovakia, Hungary and Greece (Fauna Europaea 2013) and Spain (Camargo 1992) should be substantiated. According to Vladimír Košel's personal communication, *E. monostrata* does not occur in Slovakia and the Czech Republic. The southernmost recent record has been made in Bulgaria (Moskova & Uzunov 2011). The species was also recorded from the Upper Tisza (Zakarpattia, Ukraine) (Afanasyev et al. 2013). The two latter records were made from mountain rivers that are not typical habitats of *E. monostrata*, which prefers lowland lakes and less often rivers (Agapow & Bielecki 1992, Jueg 2013). Moreover, the records are not based on morphological examinations. Therefore, the southern frontier of the species range remains unclear. *Erpobdella monostrata* has not been found in the vast interspace between its localities in Central Europe and the new locality in the Voronezh Oblast (Fig. 3). It is still unknown whether *E. monostrata* occurs in Ukraine but we may hypothesize that the species could be either overlooked or confused with related erpobdellid leeches between its eastern and western localities recorded in the previous studies. This species might be recorded as *Erpobdella testacea*

by Lukin (1929) for the Kharkiv Oblast of north-eastern Ukraine.

There are no impermeable geographical barriers in Eastern Europe that could disrupt the range of *E. monostrata*. However, our analysis revealed a substantial genetic differentiation between western and eastern populations of the species. These differences can be caused by isolation by distance or distinct postglacial colonization histories that have been already reported for freshwater leeches in the Western Palaearctic (Trontelj & Utevsky 2012).

The lack of Eastern European records may be also explained by low population sizes and the scarcity of suitable habitats. According to previous records, in 2000 only one individual of this species designated as "*Erpobdellidae* sp." was found in the Sinyutino bog situated in the Usman Pine Forest (Prokin & Silina 2007). It might be supposed that recent climate changes, i. e. exceptionally warm and dry springs and summers in 2010 and 2011 (All-Russian Research Institute of Hydrometeorological Information 2013) and forest wildfires (see above) have favoured a population expansion in the Ugolnoe Lake and after that *E. monostrata* could be found during an ecological survey. Moreover, we cannot suppose any invasive pathways and vectors for this species' range expansion. We can further speculate that the environmental changes are not favourable for *E. monostrata* in the long term as this leech prefers colder boreal freshwaters of Europe.

This research suggests that *E. monostrata* occurs from the Netherlands to the European Russian Federation. Its range may extend even more easterly where favourable habitats and climate conditions occur.

Acknowledgements

This research was supported in part by a grant from the Slovenian-Ukrainian Intergovernmental Science & Technology Cooperation Program. The pictures and DNA sequences of *E. monostrata* and *E. vilnensis* were obtained using the facilities and equipment of the Department of Biology, University of Ljubljana, courtesy of Prof. Boris Sket and Prof. Peter Trontelj. Our special thanks go to Andrei Ateasov for mapping occurrence records of *E. monostrata*. We are very thankful to Uwe Jueg and Vladimír Košel for the important information on the geographical distribution of *E. monostrata*.

References

- Afanasyev, S. A., Lietitskaya, Y. N. & Manturova, O. V. 2013. Altitude distribution and structural organization of hydrobionts' communities in the rivers of the mountainous part of the Tisa River basin. *Hydrobiological Journal* 49(4): 16–25.

- Agapow, L. & Bielecki, A. 1992. A systematic study on the complex species *Erpobdella testacea* (Savigny, 1820) (Hirudinea, Erpobdellidae). Genus 3(4): 185-199.
- & Nabodnik, J. 2006. Fauna of leeches (Hirudinea) in postglacial tarns in the Dobiegniewskie Lakeland. TEKA Komisji Ochrony i Kształtowania Środowiska [Teki Kom. Ochr. Kszt. Środ. Przyr.] 3: 13-18.
- , Korościński, B. & Nadobnik, J. 2008. Preliminary studies on the fauna of Hirudinea and Mollusca in Lake Lubikowskie. Limnological Papers 3: 9-18.
- , Nadobnik, J. & Korościński, B. 2006. Invertebrates fauna of the Lubiatowskie lakes in the north-west part of the Noteć primeval forest. Acta Agrophysica 7(2): 289-296.
- All-Russian Research Institute of Hydrometeorological Information – World Data Center. World Wide Web electronic publication. www.meteo.ru [accessed 17-Sept-2013].
- Apakupakul, K., Siddall, M. E. & Bureson, E. M. 1999. Higher level relationships of leeches (Annelida: Clitellata: Euhirudinea) based on morphology and gene sequences. Molecular Phylogenetics and Evolution 12: 350-359.
- Bielecki, A., Cichocka, J., Jeleń, I., Świątek, P. & Adamiak-Brud, Ż. 2011a. A checklist of leech species from Poland. Wiadomości Parazytologiczne 57(1): 11-20.
- , Jawniak, A. & Kalinowska, J. 2004. The leeches (Hirudinea) of the "Karas Lake" reserve in Poland. Lauterbornia 52: 33-38.
- , Świątek, P., Cichocka, J., Ropelewska E., Jeleń, I. & Adamiak-Brud, Ż. 2011b. Pijawki (Hirudinida) wód powierzchniowych Olsztyna. Forum Faunistyczne 1(1): 12-34.
- Birky, C. W. Jr. 2013. Species detection and identification in sexual organisms using population genetic theory and DNA sequences. PLoS ONE 8(1): e52544.
- Bitner, K. B., Mullen, G. & Mullen, K. 2001. Review of wildfire effects on chemical water quality. Los Alamos National Laboratory publication No. LA-13826-MS. 25 pp.
- Camargo, J. A. 1992. Macroinvertebrate responses along the recovery gradient of a regulated river (Spain) receiving an industrial effluent. Archives of Environmental Contamination and Toxicology 23: 324-332.
- Enting, K. & Arndt-Dietrich, I. 2012. Records of rare invertebrates from waters in the lowlands of North Rhine-Westphalia (Clitellata: Hirudinea; Insecta: Ephemeroptera, Coleoptera) (North West Germany). Lauterbornia 74: 81-90.
- Fauna Europaea 2013. World Wide Web electronic publication. http://www.faunaeur.org/distribution_table.php [accessed 05-Jan-2014]
- Fleituch, T. 2003. Structure and functional organization of benthic invertebrates in a regulated stream. International Review of Hydrobiology 88(3-4): 332-344.
- Folmer, O. M., Black, M., Hoeh, R., Lutz, R. & Vrijenhoek, R. 1994. DNA primers for amplification of mitochondrial cytochrome C oxidase subunit I from diverse metazoan invertebrates. Molecular Marine Biology & Biotechnology 5: 304-313.
- Grosser, C. 1996. Egelfauna des Biosphärenreservats "Mittlere Elbe" bei Dessau, Sachsen-Anhalt (Hirudinea). Lauterbornia 26: 95-98.
- 2003. Erstnachweis von *Dina apathyi* (Hirudinea: Erpobdellidae) in Deutschland. Lauterbornia 47: 59-63.
- Haaren, T., van, Hop, P., Soes, M. & Tempelman, D. 2004. The freshwater leeches (Hirudinea) of The Netherlands. Lauterbornia 52: 113-131.
- Haesloop, U. 2002. Bemerkenswerte Egelfunde aus Gräben eines nordwestdeutschen Flußmarschgebietes (NSG "Westliches Hollerland", Bremen). Lauterbornia 44: 29-36.
- Hall, T. A. 1999. BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. Nucleic Acids Symposium Series 41: 95-98.
- Jueg, U. 1998. Bemerkenswerte Egel (Hirudinea) und Krebssegel (Branchiobdellida) in Mecklenburg-Vorpommern. Lauterbornia 32: 29-47.
- 2013. Rote Liste der gefährdeten Egel und Krebssegel Mecklenburg-Vorpommerns, 1. Fassung. 56 pp., Ministerium für Landwirtschaft, Umwelt und Verbraucherschutz Mecklenburg-Vorpommern.
- , Grosser, C. & Pešić, V. 2013. Notes on the leech fauna (Hirudinea) of Kyrgyzstan. Lauterbornia 76: 103-109.
- , Schurig, B. & Sluschny, H. 2007. Beitrag zur Flora und Fauna des Neustädter Sees (Mecklenburg-Vorpommern) – Pflanzen, Pilze, Mollusken und Egel. Mitteilungen der NGM 7(1): 1-21.
- Khlyzova, N. Y., Prokin, A. A., Starodubtseva, E. A., Govorov, V. V. & Tkachenko, A. V. 2007. Materials for the study of terraced ponds of the Usman and Khrenovoe Pine Forests (I): Distribution, origin, anthropogenic transformation, typology, recurrence of the hydrological regime. Trudy Voronezhskogo Gosudarstvennogo Zapovednika 24: 234-289. [In Russian]
- Koperski, P. 2010. Urban environments as habitats for rare aquatic species: the case of leeches (Euhirudinea, Clitellata) in Warsaw freshwaters. Limnologica 40: 233-240.
- Koszalka, J. 2012. Effect of environmental factors on communities of bottom fauna in littoral zones of ten lakes in the Wel River catchment. Polish Journal of Environmental Studies 21(5): 1273-1278.
- Lukin, E. I. 1929. Biological notes on the leeches of the basin of the Donets River. Trudy Kharkivskogo Tovarystva Doslidnykiv Pryrody 52(1): 33-76. [In Russian]
- 1962. Leeches (Fauna of Ukraine). 196 pp., Kyiv (Vydavnytstvo AN URSR). [In Ukrainian]
- 1976. Leeches of fresh and saline waters (Fauna of the USSR. Leeches). 484 pp., Leningrad (Nauka). [In Russian]
- Meyer, G. A., Wells, S. G. & Jull, A. J. T. 1995. Fire and alluvial chronology in Yellowstone National Park: climatic and intrinsic controls on Holocene geomorphic processes. Geological Society of America Bulletin 107(10): 1211-1230.

- Moskova, G. D. & Uzunov, Y. I. 2011. The macrozoobenthos of the Rilska River, southwest Bulgaria. *Annuaire de l'Universite de Sofia "St. Kliment Ohridski"*. Faculte de Biologie, Livre 1 – Zoologie. Tome 99: 53–70.
- Nabodnik, J., Agapov, L. & Korościński, B. 2004. The importance of the "Santockie Zakole" nature reserve for preservation of biological diversity and tourism. *TEKA Komisji Ochrony i Kształtowania Środowiska [Tekę Kom. Ochr. Kszt. Środ. Przyr.]* 1: 157–161.
- Nakano, T. 2012. A new species of *Orobdeella* (Hirudinida, Arhynchobdellida, Gastrostomobdellidae) and redescription of *O. kawakatsurum* from Hokkaido, Japan with the phylogenetic position of the new species. *ZooKeys* 169: 9–30.
- Nesemann, H. & Moog, O. 2002. Hirudinea. – Part III. 9 pp. in: Moog, O. (ed.) *Fauna Aquatica Austriaca*, Edition 2002. Vienna (Wasserwirtschaftskataster, Bundesministerium für Land und Forstwirtschaft, Umwelt und Wasserwirtschaft).
- & Neubert, E. 1999. Annelida, Clitellata. Branchiobdellida, Acanthobdellea, Hirudinea. – Süßwasserfauna von Mitteleuropa 6/2. 178 pp., Heidelberg (Spektrum Akad. Verlag).
- Oceguera-Figueroa, A., Phillips, A. J., Pacheco-Chaves, B., Reeves, W. K. & Siddall, M. E. 2011. Phylogeny of macrophagous leeches (Hirudinea, Clitellata) based on molecular data and evaluation of the barcoding locus. *Zoologica Scripta* 40(2): 194–203.
- Pfeiffer, I., Brenig, B. & Kutschera, U. 2005. Molecular phylogeny of selected predaceous leeches with reference to the evolution of body size and terrestriality. *Theory in Biosciences* 124: 55–64.
- Prokin, A. A. & Silina, A. E. 2007. Materials for the study of terraced ponds of the Usman Pine Forest (III): Macrozoobenthos. *Trudy Voronezhskogo Gosudarstvennogo Zapovednika* 25: 300–367. [In Russian]
- Raczyńska, M. & Chojnacki, J. 2009. The structure of macrozoobenthic communities in the Tywa River, a right-bank tributary of the Oder River (northwest Poland). *International Journal of Oceanography and Hydrobiology* 38(3): 31–42.
- Ronquist, F. & Huelsenbeck, J. P. 2003. MrBayes 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics* 19: 1572–1574.
- Schenkova, J., Jarkovský, J. & Helešic, J. 2007. Strategies of coexistence of two species: *Erpobdella octoculata* and *E. vilnensis* (Hirudinea: Erpobdellidae). *International Review of Hydrobiology* 92(4–5): 527–538.
- Siddall, M. 2002. Phylogeny of the leech family Erpobdellidae (Hirudinida: Oligochaeta). *Invertebrate Systematics* 16: 1–6.
- Strauss, E. A. & Lamberti, G. A. 2000. Regulation of nitrification in aquatic sediments by organic carbon. *Limnology and Oceanography* 45: 1854–1859.
- Szczęsny, B. 2005. Some groups of benthic invertebrates and the physico-chemical conditions in the streams of the Magurski National Park in the Beskid Niski Mts (Northern Carpathians). *Nature Conservation* 61: 9–27.
- Tamura, K., Peterson, D., Peterson, N., Stecher, G., Nei, M. & Kumar, S. 2011. MEGA5: Molecular evolutionary genetics using maximum likelihood, evolutionary distance, and maximum parsimony. *Molecular Biology and Evolution* 28: 2731–2739.
- Tanabe, A. S. 2011. Kakusan4 and Aminosan: two programs for comparing nonpartitioned, proportional and separate models for combined molecular phylogenetic analyses of multilocus sequence data. *Molecular Ecology Resources* 11: 914–921.
- Trajanovski, S., Albrecht, C., Schreiber, K., Schultheiß, R., Stadler, T., Benke, M. & Wilke, T. 2010. Testing the spatial and temporal framework of speciation in an ancient lake species flock: the leech genus *Dina* (Hirudinea: Erpobdellidae) in Lake Ohrid. *Biogeosciences Discussions* 7: 5011–5045.
- Trontelj, P. & Utevsky, S. Y. 2012. Phylogeny and phylogeography of medicinal leeches (genus *Hirudo*): fast dispersal and shallow genetic structure. *Molecular Phylogenetics and Evolution* 63(2): 475–485.
- , Sket, B., Dovč, P. & Steinbrück, G. 1996. Phylogenetic relationships in European erpobdellid leeches (Hirudinea: Erpobdellidae) inferred from restriction-site data of the 18S ribosomal gene and ITS2 region. *Journal of Zoological Systematics and Evolutionary Research* 34: 85–93.
- Utevsky, S., Zigmajster, M., Atevasov, A., Zinenko, O., Utevska, O., Utevsky, A. & Trontelj, P. 2010. Distribution and status of medicinal leeches (genus *Hirudo*) in the Western Palaearctic: anthropogenic, ecological, or historical effects? *Aquatic Conservation: Marine and Freshwater Ecosystems* 20: 198–210.
- Utevsky, S. Y., Son, M. O., Dyadichko, V. G. & Kaygorodova, I. A. 2012. New information on the geographical distribution of *Erpobdella vilnensis* (Liskiewicz, 1925) (Hirudinida, Erpobdellidae) in Ukraine. *Lauterbornia* 75: 75–78.
- Westendorff, M., Kalettka, T. & Jueg, U. 2008. Occurrence of leeches (Hirudinea) in different types of water bodies in northeast Germany (Brandenburg). *Lauterbornia* 65: 153–162.
- Zhivotova, E. N. & Koroteeva, O. A. 2002. To the study of the hydrochemical regime of some waters of the Usman Pine Forest. Pp. 221–228 in: *Hydrobiological studies of some reservoirs of the Central Russian forest-steppe*. Voronezh.