

Scientific note

## Is the Sardinian grass snake (*Natrix natrix cetti*) an active hunter in underground environments?

(Serpentes, Colubridae)

Enrico Lunghi, Cristian Mascia, Manuela Mulargia & Claudia Corti

*Natrix natrix cetti* is endemic to Sardinia (Italy) (Vanni & Cimmaruta 2010). Information on the biology of this species is scarce. On several occasions (January 2016, September 2016 and twice in May 2017) we documented the underground activity of *N. natrix cetti* (Fig. 1). Snakes were observed in four caves located in four different Sardinian districts: Monte Albo (15 m from cave entrance), Supramonte (5.4 m), Sulcis-Inglesiente (75 m), Ogliastra (12 m). Caves were sub-horizontal and thus, snakes' presence was not considered accidental (Culver & Pipan 2009). *N. natrix cetti* is supposed to be one of the potential predators of the Sardinian *Hydromantes*, a group of terrestrial salamanders able to inhabit all year round underground environments (Lanza et al. 2006, Lunghi et al. 2015). Indeed, individuals of *N. natrix cetti* were found in caves inhabited by Sardinian cave salamanders (Lanza et al. 2006) where they likely entered looking for prey.

Cave predators may play an important role in underground ecosystems, being one of the major carriers of allochthonous organic matter (Culver & Pipan 2009). Understanding the role of *N. natrix cetti* in the underground food web may add useful information for the conservation of hypogean ecosystems (Culver & Pipan 2009), along with data on the dietary habits of this critically endangered snake. Furthermore, in case *Hydromantes* turned out to be a significant part of the diet of *N. natrix cetti*, it is of great importance to investigate the relationship occurring between *N. natrix cetti* and the threatened cave salamanders.

**Acknowledgements.** E. Lunghi was supported by The Mohamed bin Zayed Species Conservation Fund. The study was authorized by Regione Autonoma della Sardegna (prot. 4749 of 7 March, 2017).



Fig. 1. *Natrix natrix cetti* ♀ found in a horizontal mine 75 m far from the entrance.

### References

- Culver, D. C. & Pipan, T. (eds) 2009. The biology of caves and other subterranean habitats. 254 pp., New York (Oxford University Press).
- Lanza, B., Pastorelli, C., Laghi, P. & Cimmaruta, R. 2006. A review of systematics, taxonomy, genetics, biogeography and natural history of the genus *Speleomantes* Dubois, 1984 (Amphibia Caudata Plethodontidae). *Atti del Museo Civico di Storia Naturale di Trieste* 52: 5–135.
- Lunghi, E., Manenti, R. & Ficetola, G. F. 2015. Seasonal variation in microhabitat of salamanders: environmental variation or shift of habitat selection? *PeerJ* 3: e1122.
- Vanni, S. & Cimmaruta, R. 2010. *Natrix cetti* Gené, 1839. Pp. 538–545 in: Corti, C., Capula, M., Luiselli, L., Razzetti, E. & Sindaco, R. (eds). *Fauna d'Italia*, vol. 45 Reptilia. Bologna (Edizioni Calderini).

---

Enrico Lunghi (corresponding author), Department of Biogeography, University of Trier, Trier, Germany; and Natural History Museum of University of Florence, Zoological Section "La Specola", Firenze, Italy; and Natural Oasis, Prato, Italy; e-mail: enrico.arti@gmail.com

Cristian Mascia, Sardegna Foreste – Fo.Re.S.T.A.S., Gairo, Italy  
Manuela Mulargia, CEAS Santa Lucia, Siniscola, Italy

Claudia Corti, Natural History Museum of University of Florence, Zoological Section "La Specola", Firenze, Italy