Verlag Dr. Friedrich Pfeil ISSN 0936-9902

Ichthyological Exploration of Freshwaters

An international journal for field-orientated ichthyology

Volume 30 Number 4



Ichthyological Exploration of Freshwaters An international journal for field-orientated ichthyology

Volume 30 · Number 4 · July 2021

pages 289-384, 42 figs., 13 tabs.

Managing Editor

Paulo H. F. Lucinda, Laboratório de Ictiologia Sistemática, Universidade Federal do Tocantins, P.O. Box 136, 77500-000 Porto Nacional, TO, Brazil E-mail lucinda@uft.edu.br

Honorary Editor

Maurice Kottelat, Rue des Rauraques 6, CH-2800 Delémont, Switzerland

Associate Editors

Ralf Britz,	Senckenberg Natural History Collections Dresden, Museum of Zoology, Dresden, Germany
Kevin W. Conway,	Department of Wildlife and Fisheries Sciences, Texas A&M University, College Station, USA
Jörg Freyhof,	Leibniz-Institute of Freshwater Ecology and Inland Fisheries, Berlin, Germany
Sven O. Kullander,	Naturhistoriska Riksmuseet, Stockholm, Sweden
Heok Hee Ng,	Lee Kong Chian Natural History Museum, National University of Singapore, Singapore

Editorial Advisory Board

Museum and Art Gallery of the Northern Territory, Darwin, Australia
Naturhistorisches Museum, Bern, Switzerland
Museu de Zoologia, Unicamp, Campinas, Brazil
South African Institute for Aquatic Biodiversity, Grahamstown, South Africa
Lee Kong Chian Natural History Museum, National University of Singapore, Singapore

Ichthyological Exploration of Freshwaters is published quarterly

Subscriptions should be addressed to the Publisher:

Verlag Dr. Friedrich Pfeil, Wolfratshauser Str. 27, 81379 München, Germany PERSONAL SUBSCRIPTION: EURO 100 per Year/volume – 4 issues (includes surface mail shipping) INSTITUTIONAL SUBSCRIPTION: EURO 180 per Year/volume – 4 issues (includes surface mail shipping)

> Manuscripts should be addressed to the Managing Editor: Paulo H. F. Lucinda, lucinda@uft.edu.br

> > CIP-Titelaufnahme der Deutschen Bibliothek

Ichthyological exploration of freshwaters : an international journal for field-orientated ichthyology. – München : Pfeil. Erscheint jährl. viermal. – Aufnahme nach Vol. 1, No. 1 (1990) ISSN 0936-9902

Vol. 1, No. 1 (1990) -

Copyright © 2021 by Verlag Dr. Friedrich Pfeil, München, Germany All rights reserved.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted in any form or by any means, electronic, mechanical, photocopying or otherwise, without the prior permission of the copyright owner. Applications for such permission, with a statement of the purpose and extent of the reproduction, should be addressed to the Publisher, Verlag Dr. Friedrich Pfeil, Wolfratshauser Str. 27, 81379 München, Germany.

Printed by PBtisk a.s., Příbram I - Balonka

ISSN 0936-9902 Printed in the European Union

Verlag Dr. Friedrich Pfeil, Wolfratshauser Str. 27, 81379 München, Germany Phone + 49 89 5528600-0 · Fax + 49 89 5528600-4 · E-mail: info@pfeil-verlag.de · www.pfeil-verlag.de Ichthyol. Explor. Freshwaters, Vol. 30, No. 4, pp. 377–384, 5 figs., 3 tabs., July 2021 © 2021 by Verlag Dr. Friedrich Pfeil, München, Germany – ISSN 0936-9902 LSID: http://zoobank.org/urn:lsid:zoobank.org:pub:3D2AC8FC-D4BC-4430-BAD5-F56074F5FCBC DOI: http://doi.org/10.23788/IEF-1167 Published 24 June 2021

The first record from Laos of *Plotosus canius* (Teleostei: Plotosidae)

Kent G. Hortle* and Somphone Phommanivong**

We record the first capture from Laos of the diadromous eel-tail catfish *Plotosus canius* as a single specimen of total length 1111 mm from Phapheng Channel, which is the most easterly anabranch of the Mekong in the Khone Falls system in southern Laos. The capture location is about 2.5 km downstream of Phapheng Waterfall and about 2.1 km upstream of the Cambodian border with Laos. As the location is about 720 km from the sea and at least 500 km upstream of any possible saline influence, this record shows that some large individuals of this diadromous species, which breeds in brackish waters, may penetrate long distances into freshwaters, where they feed on crabs and molluscs.

Introduction

Plotosus canius is an eel-tail catfish species widespread in the Indo-pacific Region from Sri Lanka to New Guinea. It is found in estuaries and turbid coastal waters (Gomon, 1984), and in the 'lower parts' of rivers, including the Mekong River system in Cambodia and Vietnam (Rainboth, 1996); however the upstream limit of its distribution in the Mekong or other rivers has not been defined.

Plotosus canius is important in coastal and estuarine fisheries (Leh et al., 2012; Usman et al., 2013a). It has an extended breeding season (Usman et al., 2013a–b) and likely spawns in brackish water, based on optimal sperm survival at 0.9 % salinity (Amornsakun et al., 2018). *Plotosus canius* produces relatively few large eggs, consistent with parental care (Usman et al., 2013a; Amornsakun et al., 2018). It is possible that male fish construct

nests and guard their fry, as described for another estuarine plotosid *Cnidoglanis macrocephalus* (Laurenson et al., 1993) and for a freshwater plotosid *Tandanus tandanus* (Whitley, 1941).

Kottelat (2001) recorded 415 indigenous species of fish from the Mekong River basin in Laos. Among these he included *Plotosus canius* as being: 'tentatively recorded from Laos; its presence needs confirmation'. This tentative record by Kottelat referred to a note by Roberts (1993) who listed fishes observed or possibly known to fishermen at Hang Khone Village on the southern (downstream) shore of Khone Island, downstream of Khone Falls and about 800 m across the river from the border with Cambodia (Figs. 1–2). Roberts' species list for Khone Falls reads as follows: "?*Plotosus canius* (not observed)", as it was based only on interviews with fishermen.

^{*} Charles Sturt University, Institute for Land, Water and Society, 386 Elizabeth Mitchell Drive, Thurgoona NSW 2640, Australia. E-mail: khortle@csu.edu.au

^{**} Don Sahong Power Company, Lao-Thai Friendship Avenue, 374/15 Vat Nak Village, Sisatthanak District 0104, Vientiane, Laos. E-mail: somphone05@gmail.com



Fig. 1. Mekong River basin showing the location of Khone Falls. KF, Khone Falls; St, Stung Treng; Kr, Kratie; PP, Phnom Penh; Vt, Vientiane. Black lines are national borders; the red line is the basin boundary.

Khone Falls refers to the location where the Mekong River splits into seven main anabranches which flow southwards across the Great Fault Line of southern Laos, dropping about 20–30 m over about 5 km (Figs. 1–2). The fast-flowing channels of Khone Falls are punctuated by rapids, cascades and waterfalls which form partial or complete barriers to upstream migration by fish, depending upon species, size, location and season (Roberts & Baird, 1995).

Methods

From November 2014 to October 2020 we monitored fish abundance and diversity at Khone Falls, using several kinds of gillnets, traps (including fyke nets) and cast-nets at 12 locations upstream and downstream of Khone Falls (Fig. 2). Gillnets and traps are passive fixed gears which were set overnight, beginning one hour before sunset. Cast-netting began about one hour after sunrise for about 30 minutes, after which the gillnets and traps (including fyke nets) were retrieved. All fish were identified then measured and weighed individually. More details are provided in Hortle et al. (2014) and in other reports in preparation. The monitoring program is ongoing, supported by 16 fishermen who are paid to work full-time for the Don Sahong Hydropower Company to monitor fish and fisheries, to carry out works to improve fish passage, and to assist in fisheries management (Hortle & Phommanivong, 2019). In October 2020, the fishermen were aged 25-57 years; they had all fished since childhood and they had 16-48 years of fishing experience (total 483 man-years).

Results

From November 2014 to October 2020, the 16 fishermen caught approximately 260000 fish in 179 species in the monitoring program. Among these fish, only a single Plotosus canius was captured. The fish was caught in a fyke net set overnight (5-6 August 2016) in about 1 m of water near the eastern shore of Phapheng Channel, the most easterly of the Khone Falls channels, as shown in Figure 2. Capture coordinates are 13°56.4' N 105°59.1' E. The location is about 2.5 km downstream of Phapheng Falls and about 2 km upstream of the Cambodian border. The channel is about 300 m wide and more than 20 m deep in places, with a bedrock and sand substrate, and the river is generally fast-flowing and turbulent. During August 2016, the river was in flood and was highly turbid, as is usual for the monsoon season (Fig. 3).

The *Plotosus canius* specimen was alive when caught. It was euthanized at the site laboratory and measured 1111 mm total length (TL) and 1060 mm standard length (SL) and it weighed 8200 g (± 20 g) (Fig. 4). The fish was readily identifiable by key characters as noted by Gomon (1984) and Rainboth (1996). Diagnostic of Plotosidae, its second dorsal, caudal and anal fins were confluent, and it possessed a dendritic organ. *Plotosus* is distinguished from the only other Mekong plotosid genus, *Cnidoglanis*, by having gill membranes which are free from the isthmus,



Fig. 2. Khone Falls showing the main Mekong anabranches and the capture location of *Plotosus canius*. Red lines indicate rapids, cascades or waterfalls. DS, Don Sahong Dam Site. Arrows indicate flow direction, northwest to southeast.



Fig. 3. Phapheng Channel (right) and Sadam Channel (left) looking upstream towards Phapheng Falls on 6 August 2016. The capture location is in the right foreground, near the edge of Phapheng Channel.

and a second dorsal fin which originates above or behind the pelvic-fin insertion, as evident in Figure 4. *Plotosus canius* is distinguished from the only other Mekong congener, *P. lineatus*, by a pair of long nasal barbels which extend beyond the eye, and by a uniformly coloured body, as evident in Figures 4 and 5.

We determined that the fish was male by the size and shape of its urogenital papilla and its broad head (Usman et al., 2013a; Amornsakun et al., 2018). During dissection we noted that its testes were well-developed, but not active. Its undifferentiated gut was approximately half-full of fragments of crabs and molluscs, with some detritus and sediment, with the food items dispersed along the length of its gut. It had 5 rows of large conical or molariform mandibular teeth (Fig. 5), as well as grinding maxillary, vomerine and pharyngeal teeth, as noted and illustrated by Usman et al. (2013a: fig. 4).

Plotosus canius is not regionally rare and the specimen was large, so we did not retain it, rather it was eaten by the local fishermen who rely upon their daily catches for food. We consider that the photographs of the specimen are sufficient to document its occurrence in Laos for the first time.

Discussion

Plotosus canius is apparently rarely caught at Khone Falls, given its extremely low incidence in our samples and based on its unfamiliarity to the 16 experienced full-time fishermen, none of whom had seen a specimen of this species before the capture in 2016. They all referred to it as likely to be a kind of 'pa duk', which is the Lao name for *Clarias* species (Clariidae), true freshwater catfishes which are somewhat similar in general appearance to *Plotosus*. The fishermen subsequently heard some tentative reports that over many years of fishing at Khone Falls some of their friends may have caught one or two specimens of *P. canius*, which were however much smaller in size (~50 cm TL) than the specimen recorded herein.

Gomon (1984) noted that *Plotosus canius* is 'reportedly caught around 80 cm and may reach 150 cm in length' but cited no actual records. Kottelat (2001) noted that this species reaches 'up to 850 mm SL' (approximately 890 mm TL), presumably based on his field notes and museum records. Recent literature records, from studies in coastal or estuarine habitats, include measurements of 3272 fish, among which the largest *P. canius* recorded was 700 mm TL (Table 1), so the specimen from Khone Falls may be the largest specimen ever recorded. The fish from Khone Falls was a mature male; while we could find no records of the length at maturity of males, females have been recorded as mature at 350 mm TL (Dewanti et al., 2012).

Plotosus canius is commonly caught in the Mekong River delta in Vietnam (Fig. 1), from where Vu et al. (2020) recorded its capture by trawling at three sites which covered the range of salinity along the Mekong, from coastal brackish water to completely fresh water near the border with Cambodia, about 200 km inland. Databases of catches recorded by Mekong River fishers and compiled by the Mekong River Commission from 2003 to 2013 (unpublished) also show *Plotosus canius* as a common record in the Mekong distributaries in the Vietnam delta, but it was very rarely recorded in the Mekong in Cambodia. However, *P. canius*

Table 1. Literature records of Plotosus canius. Total length (TL) in millimetres.

Source	Sex	n	TL Minimum	TL Maximum	Location
Sinha (1981)	Male	326	114	612	India
	Female	501	86	683	
Sinha (1984)	Mixed	480	38	538	India
Khan et al. (2002)	Female	52	365	692	Bangladesh
Dewanti et al. (2012)	Female	30	350	700	Indonesia
Leh et al. (2012)	Mixed	1390	41	520	Malaysia
Usman et al. (2013b)	Female	32	412	600	Malaysia
Usman et al. (2016)	Male	172	228	629	Malaysia
	Female	169	246	600	-
Amornsakun et al. (2018)	Female	120	409	605	Thailand
Total		3272	38	700	



Fig. 4. Plotosus canius, 1060 mm SL, not preserved Phapheng Channel, Khone Falls, Laos.



Fig. 5. *Plotosus canius*, 1060 mm SL, not preserved, anterior view showing its large conical or molariform mandibular teeth.

might in fact be more common in Cambodia than reported by fishers who rarely or never see it, and who may misidentify it as a species of *Clarias*.

Similarly to some other primarily marine fishes (Roberts & Baird, 1995), Khone Falls may be the upstream limit of the distribution of *Plotosus canius* in the Mekong River, as the fast-flowing anabranch channels which cross the falls are probably an impenetrable barrier to this species, which is adapted to relatively quiescent coastal and estuarine waters. The capture of this large specimen of *Plotosus canius* 720 km upriver, and at least 500 km upstream of any saline influence, suggests that this species gains some significant benefit(s) from persisting in ascent of freshwater rivers. As discussed at length by Gross (1987), marine fishes may move into freshwaters as a direct response to environmental factors such as temperature, water level and food availability, or to avoid competition. The Mekong is noteworthy for high diversity and productivity of crabs and molluscs (Cumberlidge

et al., 2012; Köhler et al., 2012; Ngor et al., 2018), which are abundant at Khone Falls where they are not heavily fished. These invertebrates likely provide a rich food source for *P. canius*, a benthic fish, which in brackish coastal waters feeds primarily on crabs and molluscs (Sinha, 1984; Leh et

Table 2. Summary of number of records of museum specimens of *Plotosus canius* by country and habitat of capture locality, from www.fishnet2.net.

Country	Marine	Unknown	Fresh-	Total
-	or	locality	water	
	brackish		habitat	
Australia	17		2	19
Bangladesh	1	1		2
Brunei	2			2
Cambodia	0	1	5	6
India	5	9	3	17
Indonesia	4	12		16
Japan	1			1
Malaysia	24	2	2	28
Myanmar	1	2	2	5
Papua New Guinea	5			5
Philippines	16			16
Singapore	12	2		14
Sri Lanka	0	4		4
Thailand	31	7	2	40
Vietnam	2	3	1	6
Unknown	0	14		14
Total	121	57	17	195

al., 2012), which we also found in the gut of this specimen caught at Khone Falls. As the fish was caught at least 500 km upstream of any marine influence, and most items in its gut were fresh, we assume that the unidentifiable fragments of molluscs and crabs originated from recent feeding in fresh water.

To compare our observation with the extent of upstream penetration by Plotosus canius in other river systems, location records of all catalogued museum specimens of P. canius were downloaded from the online portal of www.fishnet2.net, which accesses data from 76 fish collections (Table 2). The 195 records include at least 501 specimens, but 39 of the records did not provide a count of individuals. As shown in Table 2, most records were from marine or brackish water habitats, or lacked specific locality data, with only 27 records (or 9%) from freshwaters. These records are summarised in Table 3, which shows the distance from the sea to the collection locality along the main channel of the river/stream system. Apart from the six records from the Mekong system, the furthest inland record is from the Baram River in Sarawak, Malaysia, where two P. canius were collected about 260 km inland. Further research to provide a more complete understanding of the abundance in freshwater and the geographic extent of freshwater penetration by P. canius and other diadromous fishes would be of inter-

Table 3. Details of the 17 records of 20 museum specimens of *Plotosus canius* from freshwater habitats, showing distance inland of capture localities, from www.fishnet2.net.

Country	Locality	Distance inland (km)	Number of specimens	Date
Australia	Small Coastal Creek	33	1	1975
Australia	Adelaide River, Northern Territory	69	1	1996
Cambodia	Mekong River system, Tonle Sap, Chnok Trou	390	2	1962
Cambodia	Mekong River system, Tonle Sap, Chnok Trou	390	2	1962
Cambodia	Mekong River system, Tonle Sap, Chnok Trou	390	1	1962
Cambodia	Mekong River system, Tonle Sap, Chnok Trou	390	1	1962
Cambodia	Mekong River system, Tonle Sap, Chnok Trou	390	1	1962
India	Sundarbans, Uttarbhag, Lower Bengal	98	1	1940
India	Hooghly River Calcutta	130	1	1937
India	Hooghly River tributary, Calcutta	123	1	1865
Malaysia	Baram River Sarawak	260	1	1898
Malaysia	Baram River Sarawak	260	1	no data
Myanmar	Yangon	no data	2	1937
Myanmar	Yangon River system	no data	1	no data
Thailand	Chao Phraya River, Bangkok	51	1	1862
Thailand	Bang Pakong River	no data	1	2001
Vietnam	Mekong River, Can Tho	90	1	no data

est, considering the hypothesis expounded by Roberts (1989) that "fishes with diadromous life histories generally do not survive where primary freshwater fishes are dominant".

Acknowledgements

We thank the Don Sahong Power Company for supporting the monitoring program and the publication of results, and we thank Charles Sturt University for providing library access for the senior author. Chris Barlow suggested preparing this note and kindly commented on the text. We thank the 16 fishermen who worked tirelessly on the monitoring program, and especially Khamkeng Chanbualee for his efforts in fishing the deep and sometimes hazardous waters of Phapheng Channel, where he captured the single *Plotosus canius* recorded in this paper.

Literature cited

- Amornsakun, T., B. Krisornpornsan, P. Jirasatian, T. Pholrat, T. M. Pau & A. B. Hassan. 2018. Some reproductive biological aspects of gray-eel catfish, *Plotosus canius* Hamilton, 1822 spawner in Pattani Bay, Thailand. Songklanakarin Journal of Science and Technology, 40: 384–389.
- Cumberlidge, N., P. K. L. Ng & D. C. J. Yeo. 2012. Freshwater crabs of the Indo-Burma hotspot: diversity, distribution, and conservation. Pp. 102–113 in: D. J. Allen et al. (eds.), The status and distribution of freshwater biodiversity in Indo-Burma. International Union for Conservation of Nature and Natural Resources, Cambridge, UK & Gland, Switzerland.
- Dewanti, Y. R., Irwani & S. Rejeki. 2012. Studi reproduksi dan morfometri ikan sembilang *Plotosus canius* betina yang didaratkan di pengepul wilayah Krobokan Semarang. Journal of Marine Research, 12: 135–144.
- Gomon, J. R. 1984. Plotosidae [unpaged] in: W. Fischer & G. Bianchi (eds.), FAO species identification sheets for fishery purposes. Western Indian Ocean fishing area 51. Volume 3. Food and Agriculture Organisation, Rome.
- Gross, M. R. 1987. Evolution of diadromy in fishes. American Fisheries Society Symposium, 1: 14–25.
- Hortle, K. G., P. R. Hawkins, S. Phommanivong & Y. Singsua. 2014. Khone Falls fishery monitoring methods. Don Sahong Power Company, Vientiane, Laos, 55 pp. Available from dshpp.com.
- Hortle, K. G. & S. Phommanivong. 2019. Don Sahong Fisheries Management Committee activities during construction of the Don Sahong Hydropower Project, southern Lao PDR, 2016 to mid-2019. Don Sahong Power Company, Vientiane, Laos, 57 pp. Available from dshpp.com.

- Khan, M.S.A., M. J. Alam, S. Rheman, S. Mondol & M. M. Rahman. 2002. Study on the fecundity and GSI of brackish water catfish *Plotosus canius* (Hamilton-Buchanan). Journal of Biological Sciences, 2: 232–234.
- Köhler, F., M. Seddon, A. E. Bogan, D. V. Tu, P. Sri-Aroon & D. Allen. 2012. The status and distribution of freshwater molluscs of the Indo-Burma region. Pp. 66–89 in: D. J. Allen et al. (eds.), The status and distribution of freshwater biodiversity in Indo-Burma. International Union for Conservation of Nature and Natural Resources, Cambridge, UK & Gland, Switzerland.
- Kottelat, M. 2001. Fishes of Laos. Wildlife Heritage Trust, Colombo, 198 pp.
- Laurenson, L. J. B., F. J. Neira & I. C. Potter. 1993. Reproductive biology and larval morphology of the marine plotosid *Cnidoglanis macrocephalus* (Teleostei) in a seasonally closed Australian estuary. Hydrobiologia, 268: 179–192.
- Leh, M. U. C., A. Sasekumar & L.-L. Chew. 2012. Feeding biology of eel catfish *Plotosus canius* Hamilton in a Malaysian mangrove estuary and mudflat. The Raffles Bulletin of Zoology, 60: 551–557.
- Ngor, P. B., R. Sor, L. H. Prak, N. So, Z. S. Hogan & S. Lek. 2018. Mollusc fisheries and length-weight relationship in Tonle Sap flood pulse system, Cambodia. Annales de Limnologie – International Journal of Limnology, 54: 1–10.
- Rainboth, W. J. 1996. Fishes of the Cambodian Mekong. FAO, Rome, 265 pp. plus plates.
- Roberts, T. R. 1989. The freshwater fishes of western Borneo (Kalimantan Barat, Indonesia). Memoirs of the California Academy of Sciences, 14: 1–210.
- Roberts, T. R. 1993. Artisanal fisheries and fish ecology below the great waterfalls of the Mekong River in Southern Laos. Natural History Bulletin of the Siam Society, 41: 31–62.
- Roberts, T. R. & I. G. Baird. 1995. Traditional fisheries and fish ecology on the Mekong River at Khone Waterfalls in Southern Laos. Natural History Bulletin of the Siam Society, 43: 219–262.
- Sinha, M. 1981. Length-weight relationship and relative condition factor of the canine catfish-eel *Plotosus canius* Hamilton. Journal of the Marine Biological Association of India, 23: 39–43.
- Sinha, M. 1984. Food preference studies of *Plotosus canius* Hamilton and its cultural suitability. Proceedings of Indian Academy of Sciences, 93: 437–443.
- Usman, B. I., S. M. N. Amin, A. Arshad & M. A. Rahman. 2013a. Review of some biological aspects and fisheries of grey-eel catfish *Plotosus canius* (Hamilton, 1822). Asian Journal of Animal and Veterinary Advances, 8: 154–167.
- Usman, B. I., S. M. N. Amin, A. Arshad & M. A. Rahman. 2013b. Fecundity and egg size of grey-eel catfish *Plotosus canius* (Hamilton, 1822) from the coastal waters of Kampong Telok, Negeri Sembilan Peninsular Malaysia. Asian Journal of Animal and Veterinary Advances, 8: 348–354.

- Usman, B. I., S. M. N. Amin, A. Arshad & M. S. Kamarudin. 2016. Morphometric relations in the grey eel catfish *Plotosus canius* in the coastal waters of Port Dickson, Peninsular Malaysia. Journal of Environmental Biology, 37: 573–578.
- Vu, N. U., V. H. Au & P. V. Huynh. 2020. Status of fish biodiversity and fishing on Hau River, Mekong Delta, Vietnam. Annales de Limnologie – International Journal of Limnology, 56: 1–11.
- Whitley, G. P. 1941. The catfish and its kittens. The Australian Museum Magazine, 7: 306–313.

Received 16 December 2020 Revised 17 March 2021 Accepted 8 April 2021

Ichthyological Exploration of Freshwaters An international journal for field-orientated ichthyology

INSTRUCTIONS TO CONTRIBUTORS

Warning

Prospective authors should read carefully the following instructions and follow them when submitting a manuscript. Doing so significantly hastens publication and saves money and efforts. Manuscripts which do not satisfy the instructions below may be rejected at the Editor's discretion and will not be returned.

Submission of manuscripts

The original manuscript should be sent to the Editor, Paulo H. F. Lucinda, by e-mail (lucinda@uft.edu.br). Additional information is requested:

1) the name, postal and e-mail addresses, telephone and fax numbers of the corresponding author;

 the names, postal and e-mail addresses of up to four persons outside the authors' institutions who are qualified to review the paper; and

3) a statement that the material has not been published and is not considered for publication elsewhere and that it will not be submitted elsewhere unless it is rejected or withdrawn. In submitting a manuscript, the author(s) accept(s) transfer of the copyright to the Publisher.

Co-authors, corresponding author

Authors are those who have played a significant role in designing and conducting the research and in writing the manuscript. Individuals who have only collected data, provided material or financial support, or reviewed the manuscript should be listed in acknowledgments. Honorary authorship is not accepted.

Co-authors should designate a single corresponding author to whom correspondence and proofs will be sent. All correspondence regarding the paper should go through the corresponding author. Correspondence will not be sent to other co-authors and correspondence from other co-authors regarding the manuscript will neither be answered nor taken into consideration.

Format

Files. The manuscript should be submitted in DOC or RTF format only. The text, captions, tables etc. must all be included in the same file. It the manuscript includes only a few illustrations, include them in low resolution in the word file. If the manuscript includes numerous illustrations they must be submitted in a separate PDF file; send all figures in low resolution and with caption in a single file. The files should be less than 8 MB.

Text. All manuscripts are subject to editorial revision before final acceptance for publication. Nothing in the manuscript should be underlined. Titles with numerical series designations are not permitted. Titles should be brief, fewer than 20 words and should indicate clearly the field of study and the group of fishes investigated. All abbreviations should be explained in the Method section (or figure caption when appropriate) or a reference to published explanations should be provided; exceptions are very common abbreviations, such as mm, km, kg, sec, min, yr, vs., SL. Footnotes are not permitted. All measurements must be in metric units. The first page should include: title of the paper, author(s), addresses and abstract, all left justified. The text should be followed by Material Examined (if appropriate), Acknowledgments (if any), Appendix (if any) and Literature Cited, in that order. Keys are desirable in taxonomic papers. They should be dichotomous and not serially indented.

Nomenclature. Names of living organisms should follow the appropriate and current International Codes of Nomenclature. Only formal names of genera and species should be written in italics. Names of authors and publication dates of scientific names should be mentioned once, in introduction or discussion, depending where most convenient, exceptionally as a table; bibliographical references must be included in the Literature cited section. Very old and classical works can be omitted if not absolutely justified.

Language. Manuscripts should be written in English. All papers must have a concise but informative abstract in English. In taxonomic papers, the abstract must include at least clear diagnosis of the new taxa. This maybe omitted for papers including the descriptions of many new taxa; consult the editor first. A second abstract, provided by the author(s), in the language of the country or area concerned by the text is acceptable. A maximum of two abstracts is permitted.

Acknowledgments. Identify individuals by first name(s) and surname. Do not list titles, position or institution. Acknowledge individuals, not positions. Idiosyncrasy and private jokes are not permitted.

Literature cited. Format for Literature Cited is that of the most recent issue. Do not abbreviate the names of journals. For books, give full name of publishing company or institution, and city. Manuscripts in preparation, abstracts, in-house reports and other literature not obtainable through normal library channels cannot be cited. In-press manuscripts can be cited only if they have been formally accepted.

Tables. Tables should be included in the text file, at the end. Use Word format and do not anchor them. Tables must be numbered sequentially with Arabic numerals; they should have concise but self-explanatory headings. Do not insert frames, vertical rules, dotted lines or footnotes. The location of first citation of each table should be clearly indicated in the text.

Figures. Detailed instructions for the preparation of digital images are here: https://pfeil-verlag.de/en/for-authors/

For the submission of new manuscript only low resolution copies are needed. Do not send large files at this stage. Case by case, if needed, we may ask you to send the original files at the time of submission.

All maps, graphs, charts, drawings and photographs are regarded as figures and are to be numbered consecutively and in the sequence of their first citation in the text. When several charts or photographs are grouped as one figure, they must be trimmed and spaced as intended for final reproduction. Each part of such a group figure should be lettered with a lower case block letter in the lower left corner. Where needed, scale should be indicated on the figure by a scale bar.

All illustrations should be designed to fit a width of 68 or 140 mm and a depth no greater than 200 mm. Lettering should be large enough to be easily seen when reduced onto a journal column (68 mm).

If a vector-graphics program is used, the original files saved by this program and all linked files must be submitted. Do not export or save the figure in a different format (for more details see the informations on https://pfeil-verlag.de/en/for-authors/

If line drawings are scanned, the resolution must be 1200 dpi or more and the format must be bitmap (1 pixel = 1 bit).

If halftones are scanned, the resolution should never be lower than 400 dpi, applied to a width of 14 cm, even for photographs designed for column width.

Photographic prints and slides and original drawings must be scanned for submission. We will ask to send the original after acceptance of the manuscript.

Digital images should be unmodified data files as originally saved by the camera or the scanner. If the data files are modified, a copy of the original, unmodified file should be submitted too.

The decision to print in colour or in black and white any figure originally submitted in colour remains with the editor and publisher. This decision will be based on scientific justification, quality of the original, layout and other editorial, financial and production constraints. By submitting colour originals, the authors know and accept that they may be published in black and white.

Review

Each manuscript will be sent to two reviewers for confidential evaluation. When justified, the reviewer's comments will be forwarded to the corresponding author. When submitting a revised manuscript, authors should briefly indicate the reasons for disregarding any suggestion they consider unacceptable. Remember that if a reviewer had questions or did not understand you, other readers may make the same experience and the answers should be in the manuscript and not in a letter to the editor. Changes in style, format and layout requested by the Editor are nonneoptiable and non-observance will result in rejection of the manuscript.

Revised manuscripts received more than 6 months after the reviewers' comments had been sent will not be considered or will be treated as new submissions.

Proofs, Reprints and Page Charges

A PDF proof file will be sent to the corresponding author; it should be checked and returned to the Editor within one week. If corrections are not received within this delay, they may be done by the Editor, at the author's risks. Authors may be charged for any changes other than printer's error. Reprint orders must be forwarded with the corrections. The corresponding author is responsible for contacting the co-authors and forwarding their reprint orders.

The authors will receive a PDF file for personal use free of charge; high-resolution PDF files for unlimited use may be ordered. There will be no page charges and no charges for justified colour illustrations.

Ichthyological Exploration of Freshwaters

An international journal for field-orientated ichthyology

Volume 30 · Number 4 · July 2021

CONTENTS

Schneider, Nikole G., Andrew J. McKamy, Alexander M. Rubin, Heiko L. Schoenfuss, Richard W. Blob and Kelly M. Diamond: In-stream analysis of predator-prey interactions in Hawaiian stream fishes	289
Cheng, Jian-Li, Wei-Han Shao, J. Andrés López and E Zhang: <i>Tachysurus lani</i> , a new catfish species (Teleostei: Bagridae) from the Pearl River basin, South China	299
Katemo Manda, Bauchet, Jos Snoeks, Auguste Chocha Manda and Emmanuel Vreven: <i>Nan-nocharax hadros</i> (Teleostei: Distichodontidae), a new species from the Kalule Nord River in the Upemba National Park, Upper Congo basin	317
Perez, Camila F., Alexandre K. Oliveira and Júlio C. Garavello: Morphological variation of <i>Hypostomus affinis</i> (Teleostei: Loricariidae) from eastern Brazilian drainages	331
Skelton, Paul H., Jay R. Stauffer, Albert Chakona and Joshua M. Wisor: Two new species of African bubble-nesting <i>Microctenopoma</i> (Teleostei: Anabantidae) from Angola	345
Zawadzki, Cláudio Henrique and Iago de Souza Penido: <i>Hypostomus bimbai</i> and <i>H. pastin-</i> <i>hai</i> , two new depth-sheltered species of <i>Hypostomus</i> (Teleostei: Loricariidae) from the lower Rio São Francisco basin, Brazil	361
Hortle, Kent G. and Somphone Phommanivong: The first record from Laos of <i>Plotosus canius</i> (Teleostei: Plotosidae)	377

Cover photograph Hypostomus pastinhai (Photograph by Cláudio Henrique Zawadzki) Cláudio Henrique Zawadzki and Iago de Souza Penido (this volume pp. 361–375)

Articles appearing in this journal are indexed in:

AQUATIC SCIENCES and FISHERIES ABSTRACTS BIOLIS - BIOLOGISCHE LITERATUR INFORMATION SENCKENBERG CAMBRIDGE SCIENTIFIC ABSTRACTS CURRENT CONTENTS/AGRICULTURE, BIOLOGY & ENVIRONMENTAL SCIENCES and SCIE FISHLIT ZOOLOGICAL RECORD